

Competition of Orbital Antiferromagnetism With Q-Triplet-Pairing in the Ferromagnetic Superconductor UGe₂

G. Varelogiannis^a P. Thalmeier^b S. Tsonis^a

^a*Department of Physics, National Technical University of Athens, GR-15780 Athens, Greece*

^b*Max-Planck-Institute for Chemical Physics of Solids, D-01187 Dresden, Germany*

Abstract

Within a multicomponent BCS-like framework we study the coexistence and competition of various CDW and SC order parameters in the presence of a strong ferromagnetic background. We find that the competition of unconventional CDW, also called orbital antiferromagnet, with SC at finite momentum (**Q**-triplet pairing) shows unique characteristics like an extreme sensitivity on the deviations from nesting. We argue that pressure in UGe₂ creates deviations from the nesting and report a phase diagram in qualitative agreement with the observed behavior of UGe₂.

Key words: Ferromagnetism and Superconductivity, UGe₂, CDW and superconductivity

PACS: 74.20.-z

The discovery of superconductivity (SC) well inside the itinerant ferromagnetic (FM) state of UGe₂ [1] is a true challenge in the theory of superconductivity in strongly correlated systems. It appears now quite clear that SC develops only inside the FM state quite suddenly when the critical temperature T^* of a hidden order state inside the FM state is reduced towards zero by applying pressure. The existence of a true thermodynamic phase with hidden order below T^* has been suggested by specific heat experiments [2]. The FM transition is apparently of first order [3] which is very hard to reconcile with any approach that would associate superconductivity with the fluctuations of the FM or the T^* hidden order as in the quantum critical point scenario.

Within our model we obtain a pressure- temperature phase diagram that displays many of the

observed characteristics of UGe₂. We propose that T^* corresponds to a CDW formation and we have studied the competition of various CDW and SC order parameter symmetries within a BCS like mean-field framework in the presence of a fully polarized FM background, i.e. frozen spins. We use a Nambu type formalism with spinors defined by

$$\Psi_{\mathbf{k}}^\dagger = (c_{\mathbf{k}}^\dagger, c_{-\mathbf{k}}, c_{\mathbf{k}+\mathbf{Q}}^\dagger, c_{-\mathbf{k}-\mathbf{Q}}) \quad (1)$$

Nesting in the fully polarized band is responsible for the CDW transition that we associate with the T^* line in UGe₂. Pressure reduces T^* because it removes the nesting features. To discuss this phenomenon we write the electron dispersion $\xi_{\mathbf{k}}$ as a sum of particle-hole symmetric terms responsible for nesting and particle-hole asymmetric terms that represent the deviations from nesting: $\xi_{\mathbf{k}} = \gamma_{\mathbf{k}} + \delta_{\mathbf{k}}$ where $2\gamma_{\mathbf{k}} = \xi_{\mathbf{k}} - \xi_{\mathbf{k}+\mathbf{Q}}$ and $2\delta_{\mathbf{k}} = \xi_{\mathbf{k}} +$

$\xi_{\mathbf{k}+\mathbf{Q}}$. When $\delta_{\mathbf{k}} = 0$ there is particle-hole symmetry or perfect nesting at the wavevector $\mathbf{Q}$.

Our approach allows to study SC and CDW states which transform either odd or even under $\mathbf{k} \rightarrow -\mathbf{k}$ (inversion) or $\mathbf{k} \rightarrow \mathbf{k} + \mathbf{Q}$. The possible superconducting states must be odd under inversion because the spins are frozen. We therefore examine a special type of unitary triplet SC state. Our analysis shows that with these symmetry constraints we must necessarily have $\Delta_{\mathbf{k}+\mathbf{Q}}^{\mathbf{Q}} = -\Delta_{\mathbf{k}}^{\mathbf{Q}}$ where $\Delta_{\mathbf{k}}^{\mathbf{Q}}$ is the SC gap for pairing at finite momentum $\mathbf{Q}$ also called $\mathbf{Q}$ -triplet pairing or π -triplet pairing. For the zero momentum pairing state the order parameter $\Delta_{\mathbf{k}}$ must be purely imaginary. Concerning the CDW states, for states with both even and odd parity the order parameter $W_{\mathbf{k}}$ is real if $W_{\mathbf{k}} = W_{\mathbf{k}+\mathbf{Q}}$ and imaginary if $W_{\mathbf{k}} = -W_{\mathbf{k}+\mathbf{Q}}$. The CDW states which obey $W_{\mathbf{k}} = -W_{\mathbf{k}+\mathbf{Q}}$ are unconventional CDW states (UCDW) also called orbital antiferromagnetic states due to the staggered current pattern they induce on the lattice plaquettes.

A study of the coexistence and competition of all accessible SC and CDW states will be presented elsewhere. Here we focus on a single case which is perhaps the most exciting one showing unique characteristics that may plausibly correspond to the physical situation observed in the FM state of UGe_2 . It corresponds to the competition of the even parity UCDW state with the finite- $\mathbf{Q}$ SC state. To simplify the notation from now on we will note $\Delta_{\mathbf{k}}^{\mathbf{Q}} \equiv \Delta_{\mathbf{k}}$. The coupled gap equations that we obtain in that case have the following form:

$$W_{\mathbf{k}} = \sum_{\mathbf{k}'} V_{\mathbf{k},\mathbf{k}'}^{CDW} W_{\mathbf{k}'} \frac{1}{4\sqrt{\gamma_{\mathbf{k}'}^2 + W_{\mathbf{k}'}^2}} \times \left(\tanh \frac{E_+(\mathbf{k})}{2T} + \tanh \frac{E_-(\mathbf{k})}{2T} \right) \quad (2)$$

$$\Delta_{\mathbf{k}} = \sum_{\mathbf{k}'} V_{\mathbf{k},\mathbf{k}'}^{SC} \Delta_{\mathbf{k}'} \frac{1}{4\sqrt{\delta_{\mathbf{k}'}^2 + \Delta_{\mathbf{k}'}^2}} \times \left(\tanh \frac{E_+(\mathbf{k})}{2T} - \tanh \frac{E_-(\mathbf{k})}{2T} \right) \quad (3)$$

where $W_{\mathbf{k}}$ is the UCDW gap and

$$E_{\pm}(\mathbf{k}) = \sqrt{\gamma_{\mathbf{k}}^2 + W_{\mathbf{k}}^2} \pm \sqrt{\delta_{\mathbf{k}}^2 + \Delta_{\mathbf{k}}^2} \quad (4)$$

The system of gap equations (2) and (3) is very peculiar showing a unique influence of particle-hole

asymmetry on the SC versus UCDW competition. Indeed, in the UCDW gap equation (2), in the zero temperature regime, only the particle-hole symmetric terms $\gamma_{\mathbf{k}}$ of the dispersion enter. In the SC gap equation (3) instead, in the same temperature regime, the particle-hole asymmetric terms $\delta_{\mathbf{k}}$ appear. On the other hand at finite temperatures both terms enter in both gap equations.




Fig. 1. Phase diagram showing the competition of UCDW with $\mathbf{Q}$ -triplet SC as a function of the particle-hole asymmetry. The two states do not coexist but rather exclude each other.

We solve numerically the gap equations (2) and (3) considering a n.n.n tight binding model for the majority spin band of UGe_2 . In which case $\gamma_{\mathbf{k}} = t(\cos k_x + \cos k_y)$ and $\delta_{\mathbf{k}} = t' \cos k_x \cos k_y$.

We assume that the effect of pressure is to create deviations from nesting by enhancing the ratio t'/t . Some of our results are shown in Fig. 1. Bearing in mind that the relationship of pressure with t'/t is likely nonlinear, the results shown in Fig. 1 are quite similar to the pressure behaviour of UGe_2 . The UCDW gap is very rapidly reduced with particle hole asymmetry and above a critical t'/t (i.e. a critical pressure) $\mathbf{Q}$ -triplet SC suddenly appears. The fact that SC is taking its maximal critical temperature at the point where the UCDW disappears is not related to any role played by the fluctuations of the incipient UCDW order parameter since the latter is in rather strong competition with SC.

References

- [1] S.S. Saxena et al., Nature **406**, 587 (2000).
- [2] N. Tateiwa et al., Acta Phys. Pol. B 34, 515 (2003)
- [3] C. Pfeiderer and A.D. Huxley, Phys. Rev. Lett. **89**, 147005 (2002).